

Crop(s) **Barley**
Disease(s) **Powdery mildew**

Resistance management strategy for:

Group 3	Demethylation Inhibitors (DMI);
Group 5	Morpholines;
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 7 + 3	SDHI + DMI;
Group 11 + 3	Quinone outside Inhibitor (QoI) + DMI;
Group 11 + 4	QoI + Phenylamide;
Group 13	Aza-naphthalene; and
Group 11 + 7 + 3	QoI +SDHI + DMI.

Guidelines

- Fungicides should be used preventatively (as protectants), or at the very first sign of disease. If disease is established within the canopy, fungicides may not produce optimal results and there is a high chance of selection for fungicide resistance. In high disease pressure situations, integrated management approaches should be considered to reduce fungicide resistance including:
 - Removal of stubble;
 - Crop rotation (**avoid** barley on barley);
 - Control of green bridge volunteers;
 - Use of tolerant or resistant varieties; and
 - Strategic use (timing, application technique and rate) of fungicides.
- Do not** apply more than two applications per growing season of products containing any of the above listed Groups containing products. This includes in-furrow or seed treatments that have activity on powdery mildew. Combinations of in-furrow and seed treatment are counted as one application.
- Do not** apply consecutive applications of **Group 11** containing products. This includes in-furrow i.e. If a **Group 11+4** fungicide has been used in-furrow at planting, and the **Group 11** has foliar activity (as determined by label claims), the first foliar fungicide spray must not contain a **Group 11** fungicide.
- If a **Group 7** seed treatment has been used with foliar activity (as determined by label claims), the first foliar fungicide applied must not contain a **Group 7** fungicide.
- Group 7** and **Group 13** foliar fungicides must always be in a co-formulation or in mixture with a registered mixing partner with a different mode of action, with no known resistance.
- Minimise use of fungicides which are known to have compromised efficacy due to resistance.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.

