

Crop(s) Berry (generic, other than strawberry)
Disease(s) Grey mould (*Botrytis*)

Resistance management strategy for:

Group 2	Dicarboximide;
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 7 + 11	SDHI + Quinone outside inhibitors (QoI);
Group 7 + 12	SDHI + phenylpyrroles (PP);
Group 9	Anilinopyrimidine (AP);
Group 9 + 12	AP + PP;
Group 17	Hydroxylanilide;
Group 19	Chitin synthase inhibitor;

Guidelines

1. Apply a program of protectant fungicides during flowering.
2. **Do not** apply more than two consecutive sprays of **Group 2, 9, , 17, 19** or fungicides. Continue alternation of Mode of Action between successive crops.
3. Do not apply **17** (solo) as consecutive applications, or more than 3 times per season.
4. If using **Group 17** with an effective mixing partner, or a different **Group**, **do not** apply the **Group 17** containing mixtures for more than 50% of all applications.
5. If applying **Group 7** (including, **7+11** or **7+12**) fungicides, **do not** apply more than two consecutive sprays before changing to another group. **Do not** apply more than three **Group 7** sprays per season. If consecutive sprays are used, then use the same number of sprays of an alternative group before using another **Group 7**, including sprays in consecutive seasons.
6. If three or fewer sprays are applied per crop, use only one spray containing a **Group 9** or **12** containing fungicide. Use a maximum of three sprays containing **Group 9, or 12** containing fungicides, or 33% of sprays (whichever is lower).
7. **Do not** apply **Group 19** fungicides more than 6 times per season, or more than 50% of the total number of sprays.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



	Group						
	2	7 7+11 7+12	9 9+12	12 12+7 12+9	17 (solo)	17 (in mixture)	19
Maximum number of consecutive applications	2	2	2	2	1	1	2
Maximum number of sprays per crop/season	2-3	3 Or 1 in 3 (33%) whichever is lower	3 Or 1 in 3 (33%) whichever is lower	3 Or 1 in 3 (33%) whichever is lower	3 Or 1 in 3 (33%) whichever is lower	1 in 2 (50%)	6 Or 1 in 2 (50%) whichever is lower

Where there are several crop cycles per year, up to a maximum of five (5) applications containing **Group 21** can be applied.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.

