

Crop(s) **Ornamentals**
Disease(s) **Grey mould (*Botrytis*)**

Resistance management strategy for:

Group 2	Dicarboximide;
Group 9	Anilinopyrimidine (AP);
Group 9 + 12	AP + phenylpyrroles;
Group 11	Quinone outside inhibitor (QoI);
Group 11 + 3	QoI + Demethylation inhibitors (DMI); and
Group 17	Hydroxyanilide.

Guidelines

1. If three or fewer sprays are applied per season, use only one spray containing a **Group 9** fungicide (including **9+12**). If four to six sprays are applied per season, use a maximum of two sprays containing **Group 9** fungicides. If seven or more sprays are applied per season use a maximum of three sprays containing **Group 9** fungicides.
2. **Do not** apply more than two consecutive sprays of a **Group 2, 9, 11, 12 or 17** fungicides.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.

