

Crop(s) **Avocado and mango**
Disease(s) **Anthrachnose (Colletotrichum spp.)**

Resistance management strategy for:

Group 3	Demethylation inhibitors (DMI);
Group 7 + 11	Succinate dehydrogenase inhibitors (SDHI) + Quinone
	outside inhibitor (Qol); and
Group 11*	Quinone outside inhibitor (Qol).

Guidelines

19. If applying **Group 11** (including **7+11**) fungicides, **do not** apply more than two consecutive sprays before changing to another group.
20. **Do not** apply more than three **Group 11** sprays per season within the field. If consecutive sprays are used, then use the same number of sprays of an alternative group before using another **Group 11***.
21. **Do not** apply more than four **Group 3** sprays in a season. Apply no more than two consecutive sprays of a **Group 3** fungicide alone.

* *Registered for use in mangoes only.*

17

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.

