

Crop(s)	Stone fruit
Disease(s)	Blossom blight (<i>Monilinia laxa</i>), brown rot (<i>M. fructicola</i>)

Resistance management strategy for:

Group 2	Dicarboximides;
Group 3	Demethylation inhibitors (DMI);
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 7 + 11	SDHI + Quinone outside inhibitors (QoI);
Group 9	Anilinopyrimidines (AP); and
Group 11	Quinone outside inhibitors (QoI).

Guidelines

1. If applying **Group 2, 3, 7, 11** or **7+11** fungicides, **do not** apply more than two consecutive sprays of fungicides from the same group before changing to another group, this includes the last application made in-field from one season to the next.
2. **Do not** apply more than three sprays of a **Group 7, 9, 11** or **7+11** fungicide per season. Alone, tank mixed or co-formulated all count to total. If group 7 applied alone, apply **Group 7** in strict alternation with fungicides from a different cross-resistance group.
3. When mixtures are used for **Group 7** resistance management, applied as tank mix or as a co-formulated mixture, the mixture partner should provide effective disease control when used alone on the target disease and must have a different mode of action.
4. Always apply **Group 7** fungicides as per label instructions and use preventatively.
5. If consecutive sprays of fungicides from the same chemical group are applied, they must be followed by at least the same number of sprays from an alternative chemical group, before returning to the original group.

57

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



	Group					
	2	3	7	9	11	7 + 11
Maximum number of consecutive applications	2	2	2	2	2	2
Maximum number of solo sprays	2	2	2	2	2	2
Maximum number of sprays per season	2	3	3	3	3	3

6. A post-harvest foliar treatment should also be counted as an application.
7. The last blossom blight spray and the first pre-harvest brown rot spray should be regarded as consecutive applications.
8. The spray program should be considered and the resistance management strategy applied on a whole-orchard basis.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.

