

Crop(s)	Nursery e.g., vegetable seedlings, trees, ornamentals
Insect(s)	Various

Guidelines

1. Monitor regularly for insect activity. Where relevant consider the use of light, insect traps or other monitoring tools.
2. Only use insecticides when insect populations reach accepted threshold levels.
3. Always read and follow product labels and use the full recommended label rates of application. Some products place a limit on the number of times they can be applied per crop and when they can be applied.
4. Ensure the spray equipment is properly calibrated and achieves good coverage with appropriately sized spray droplets.
5. In the case of soil applied insecticides, apply once only. Apply using sufficient water that does not result in any drop point from the soil medium containing the seedling or plant. If watering is required between applications and planting, it should be done sparingly, only as required. **Avoid** water to run-through from the cells, bags etc.
6. When transplanting soil-treated seedlings or plants, ensure that the growing medium is fully transferred to the field with each seedling or plant.
7. **Rotate** between registered insecticides that have different modes of action.
8. Where possible **avoid applying consecutive applications** of insecticides that have the same mode of action within and between seasons or exceed the recommended maximum number of applications in a crop.
9. **Do not** follow a seed, seedling or soil treatment with a foliar application from the same mode of action group.
10. Time the foliar applications to the most susceptible life stage of the target pest.
11. **Do not** re-treat a spray failure with a product from the same mode of action group.
12. Avoid using insecticide tank-mixes where both active ingredients control the same insect pest as this strategy is generally not considered best practice for resistance management.
13. Practice good crop hygiene to reduce insect pressure e.g. removing severely infested seedling trays, plants or host weeds.
14. Nurseries supplying treated plants to commercial operations should clearly identify insecticides that have been applied and supply paperwork to the recipient accompanying the plant that indicates the rates and date. For treated seedlings the time of application should also be included, particularly if applied just prior to transplant in the field.

15

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Notes regarding the application of insecticides

1. Refer to the CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides.
2. There is known cross-resistance between some chemical groups e.g., **Groups 1A** and **1B**.
3. Seek advice from the manufacturers or government advisory services to determine local resistance levels for specific mode of action groups.
4. **Do not** exceed the maximum number of applications permitted on the insecticide label.
5. When using insecticides to control other pests consider the chemical group in relation to contributing to the resistance development of other insect pests.
6. When using insecticides to control insect pests consider the effect on beneficial insects and the potential to flare insect populations.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.

