

Crop(s)	Various
Insect(s)	Silverleaf whitefly (<i>Bemisia tabaci</i>)

Guidelines

1. Monitor pest numbers and apply control measures before adult populations reach high levels.
2. Select registered insecticide control measures according to the primary growth stage of the pest, the infestation level and the age and type of crop.
3. In cotton, spray decisions should be based on the Silverleaf Whitefly threshold matrix. Refer to the current [Cotton Pest Management Guide](#) for further details.
4. Where possible, utilise selective insecticides during the early stages of crop development to minimise the impact on beneficial insects.
5. Rotate between registered insecticides that have different modes of action (see table below).
6. **Do not** apply more than two consecutive applications of insecticides that have the same Mode of Action within and between seasons.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



7. The Modes of Action (groups) and registered insecticides for control of Silverleaf Whitefly are listed below:

Group*	Chemical subgroup	Active ingredient
1B	Organophosphate	acephate
3A	Pyrethroids	bifenthrin
4A	Neonicotinoids	acetamiprid, clothianidin, dinotefuran, imidacloprid, thiamethoxam
4D	Butenolides	flupyradifurone
7C	Pyriproxyfen	pyriproxyfen
9D	Pyropenes	afidopyropen
12A	Diafenthiuron	diafenthiuron
16	Buprofezin	buprofezin
23	Tetronic and Tetramic acid derivatives	spirotetramat
28	Diamides	cyantraniliprole
28 + 4A	Diamides + Neonicotinoids	cyantraniliprole + thiamethoxam
29	Flonicamid	flonicamid
36	Pyridazine pyrazolecarboxamides	dimpropyridaz
UN		clitoria ternatea extract
UNF	<i>Beauveria bassiana</i> strains	<i>Beauveria bassiana</i> strain PPRI 5339
UNM	Mineral oil	paraffinic oil

46

*Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides

8. Seek advice from the manufacturers and/or government advisory services to determine local resistance levels for particular mode of action Groups.
9. **Do not** exceed the maximum number of applications permitted on the insecticide label.
10. When using insecticides to control other pests, consider the chemical group in relation to contributing to resistance development of Silverleaf Whitefly.
11. When using insecticides to control other pests consider the effect on beneficial insects and the potential to flare Silverleaf Whitefly populations.

Notes

Not all chemical groups listed have registered products available in all crops affected by Silverleaf Whitefly. Only use products registered for use in crop to be treated.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Cultural practices

1. In vegetable crops, ensure seedlings are free of pests prior to transplanting. Inspect transplants carefully upon arrival for whitefly eggs, nymphs and adults.
2. Control alternate weed hosts of Silverleaf Whitefly 2-3 weeks before planting to reduce early population levels.
3. Clean-up crop residues:
 - a. Where moderate population levels remain after harvest, apply a registered insecticide or oil treatment effective against adults.
 - b. Plough in crops within 2-3 days of application to kill all remaining nymphs on crop foliage to reduce pest migration into new plantings.

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.

