

CONTENTS

INTRODUCTION		
Principles of resistance management		2
Effective resistance management strategies use alternations or sequences of different modes of action		2
What is resistance?		2
Resistance mechanisms		3
Mode of action, target site resistance and cross resistance		3
Alternation of chemistry		4
Use of cultural practices		4
Understanding of the insect or mite life cycle		4
Application		4
Resistance management strategy design		4
Additional information		5
SPECIFIC GUIDELINES		
CROP(S)	PEST(S)	
Banana	Banana weevil borer (<i>Cosmopolites sordidus</i>) and Rust thrips (<i>Chaetanaphothrips signipennis</i>)	6
Brassica, brassica leafy vegetables	Diamondback moth, Cabbage moth (<i>Plutella xylostella</i>)	10
Canola, Forage brassica	Diamondback moth, Cabbage moth (<i>Plutella xylostella</i>)	12
Cotton	All pests	14
Nursery	Various	15
Pasture/Winter crops	Redlegged earth mite (RLEM) (<i>Halotydeus destructor</i>)	17
Pome Fruit	Two spotted mite (<i>Tetranychus urticae</i>), European red mite (<i>Panonychus ulmi</i>)	19
Potato	Potato tuber moth / Tomato leafminer (<i>Phthorimaea operculella</i>)	21
Sorghum, Maize, Summer & Winter Grain Legumes	Heliothis/Cotton bollworm/Native budworm (<i>Helicoverpa</i> spp.)	22
Strawberries/Ornamentals	Two-spotted mite (<i>Tetranychus urticae</i>)	25
Sweet corn	Corn earworm (<i>Helicoverpa armigera</i>) aka Heliothis	27
Tomato	Heliothis/Tomato budworm (<i>Helicoverpa</i> spp.)	30
Turf	Various	33
Various	Cotton/Melon aphid (<i>Aphis gossypii</i>), Green peach aphid (<i>Myzus persicae</i>)	35
	Fall armyworm (<i>Spodoptera frugiperda</i>)	37
	Mites	41
	Silverleaf whitefly (<i>Bemisia tabaci</i>)	42
	Western flower thrips (<i>Frankliniella occidentalis</i>)	44
Wide host range of plant spp.	Serpentine leafminer (<i>Liriomyza huidobrensis</i>)	45
	Nematodes	51

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Introduction

The CropLife Australia Expert Committee on Insecticide Resistance (ECIR) has drafted insect resistance management strategies in conjunction with growers, researchers and agronomists to minimise the development of insect resistance to insecticides. These strategies provide growers with guidelines for insecticide use (and other methods) for sustainable insect control.

Principles of resistance management

Insecticide or acaricide resistance management strategies seek to minimise the selection for resistance to any one type of insecticide or acaricide. This requires an understanding of insecticides as they are grouped according to similarity of Mode of Action (MoA) in controlling insects and mites.

In practice, sequences or rotations of compounds from different MoA groups provide an effective approach to resistance management. These MoA groups are shown in the **Mode of Action Classification for Insecticides Table**.

Effective resistance management strategies use alternations or sequences of different modes of action.

The objective of Insecticide Resistance Management is to prevent or delay resistance developing to insecticides, or to help regain susceptibility in insect pest populations in which resistance has already arisen. IRM is important in maintaining the efficacy of valuable insecticides. It is usually easier to prevent resistance occurring than it is to reactively regain susceptibility.

Insecticide applications are often arranged into MoA spray windows or blocks that are defined by the stage of crop development and the biology of the pest(s) of concern. Local expert advice should always be followed with regard to spray windows and timings. Several sprays of a compound may be possible within each spray window but it is generally essential to ensure that successive generations of the pest are not treated with compounds from the same MoA group.

2

What is resistance?

Resistance to insecticides and acaricides may be defined as *'a heritable change in the sensitivity of a pest population that is reflected in the repeated failure of a product to achieve the expected level of control when used according to the label recommendation for that pest species.'*

Resistance arises through the overuse or misuse of an insecticide or acaricide against a pest species and results in the selection of resistant forms of the pest and the consequent evolution of populations that are resistant to that insecticide or acaricide.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Resistance mechanisms

There are several ways insects can become resistant to insecticidal crop protection products.

Metabolic resistance

Resistant insects may detoxify or destroy the toxin faster than susceptible insects or quickly rid their bodies of the toxic molecules. Metabolic resistance is the most common mechanism and often presents the greatest challenge. Insects use their internal enzyme systems to break down insecticides. Resistant strains may possess higher levels or more efficient forms of these enzymes. In addition to being more efficient, these enzyme systems also may have a broad spectrum of activity (i.e. they can degrade many different insecticides).

Target site resistance

The target site where the insecticide acts in the insect may be genetically modified to prevent the insecticide binding or interacting at its site of action thereby reducing or eliminating the pesticidal effect of the insecticide.

Penetration resistance

Resistant insects may absorb the toxin more slowly than susceptible insects. Penetration resistance occurs when the insect's outer cuticle develops barriers which can slow absorption of the chemicals into their bodies. This can protect insects from a wide range of insecticides. Penetration resistance is frequently present along with other forms of resistance, and reduced penetration intensifies the effects of those other mechanisms.

Behavioural resistance

Resistant insects may detect or recognize a danger and avoid the toxin. This mechanism of resistance has been reported for several classes of insecticides, including organochlorines, organophosphates, carbamates and pyrethroids. Insects may simply stop feeding if they come across certain insecticides, or leave the area where spraying occurred (for instance, they may move to the underside of a sprayed leaf, move deeper in the crop canopy or fly away from the target area).

Mode of action, target-site resistance, and cross-resistance

In the majority of cases, not only does resistance render the selecting insecticide ineffective but it often confers cross-resistance to other chemically related compounds. Compounds within a specific chemical group usually share a common target site within the pest and thus share a common Mode of Action (MoA). It is common for resistance to develop that is based on a genetic modification of this target site. When this happens, the compound loses its pesticidal efficacy. Because all compounds within the chemical sub-group share a common MoA, there is a high risk that the resistance will automatically confer cross-resistance to all the compounds in the same sub-group. It is this concept of cross-resistance within chemically related insecticides or acaricides that is the basis of the Mode of Action classification.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Alternation of chemistry

Constant use of insecticides from one chemical grouping (MoA) will increase the risk of rapid build-up of resistance to that chemical group. Alternate use of chemical groups with different MoAs will slow down the process of selection for resistance.

Use of cultural practices

Incorporation of cultural techniques for controlling an insect pest will reduce selection pressure from the insecticides. Any resistance management strategies should incorporate all available methods of control for the insect pest concerned.

Understanding of the insect or mite life cycle

A good understanding of the life cycle of the pest is essential so that control methods can be effectively targeted. An insecticide or acaricide should always be targeted at the pest growth stage that is most susceptible for that insecticide or acaricide.

Application

Label recommendation

Insecticide labels have been carefully developed to ensure the most effective control of the pest. The label should at all times be carefully read and adhered to.

4

Coverage

The majority of insecticides require good coverage of the target area to ensure the best possible chance of contact and subsequent control of the pest.

Resistance management strategy design

Crop/pest or regional strategies

The strategies below are provided on a CROP by PEST basis (e.g. Tomato - Heliothis). However, in horticultural and agricultural areas often a range of crops are grown that are attacked by a range of pests.

In many cases, a specific MoA insecticide can be used across this range of crops to control multiple pests that have the ability to move from crop to crop. There is interaction between intensive horticulture and broadacre farming, as with Diamondback Moth (DBM) in Brassica vegetables and resistance strategies that could be compromised by widespread use of insecticides for DBM control in canola.

Also, the pest complex for a specific crop will vary within production regions, especially between Northern and Southern Australia.

For this reason, CROP by PEST strategies can be flawed and further Insecticide Resistance Management (IRM) advice for specific pests should always be sought on a local basis.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



An alternative to the CROP by PEST strategy is that of “Regional strategies” such as those for Cotton, Brassicas and the Southern NSW and Northern Victorian IRM strategy for grain and annual horticultural crops.

These regional or specific crop strategies are available on the CropLife Australia website.

The overall Resistance Management Strategy of avoiding overuse of individual Modes of Action insecticides should be followed, not just on a specific crop and pest but on a broad perspective of crops and pest complex.

Additional information

Further information on Insecticide Resistance, Management Strategies and Insecticide Mode of Action can be found on the International IRAC (Insecticide Resistance Action Committee) website:

<https://www.irac-online.org/>.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Banana
Insect(s)	Banana weevil borer (<i>Cosmopolites sordidus</i>) and Banana rust thrips (<i>Chaetanaphothrips signipennis</i>)

Guidelines

1. Use only clean planting material, such as tissue culture plants or bits/suckers from a clean nursery block.
2. If re-planting into an old banana block, allow at least 6 months fallow after the old banana material has rotted down.
3. Remove weeds and trash around banana stools to allow maximum effectiveness of insecticides and to reduce sheltering sites for weevils. Application of insecticides to trash may lead to reduced control of banana weevil borer.
4. Cut up fallen and harvested pseudo-stems to reduce weevil breeding sites.
5. Monitor regularly for banana weevil borer activity by trapping (when adult weevils are active) or conduct corm damage ratings. Dying water suckers can also be an indication of banana weevil borer populations.
6. Only use insecticides when populations reach or exceed accepted threshold levels. Refer to local Department of Agriculture guidelines.
7. Only use insecticides at the registered rate of application and apply at times when the particular product will have the maximum impact, i.e., use contact insecticides only when banana weevil borer adults are active.
8. Insecticides should only be used in the years indicated in the following diagrams.
9. When applying insecticides for the control of banana weevil borer and banana rust thrips consider the impact of these products on other insects and nematodes, and vice versa.
10. For banana rust thrips control, a combination of control methods such as butt or band sprays, stem injection, bell injection and/or bunch sprays may be required.

6

Continued on next page

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



STRATEGY A: Where products other than controlled release formulations of imidacloprid are being used to control insects in bananas

MoA Group*	Chemical subgroup	Example chemical	Year 1 use	Year 2 use	Year 3 use	Year 4 use	Year 5 use	Year 6 use
1A or 1B	Carbamates Organophosphates	oxamyl ¹ or acephate ² cadusafos ¹ prothiofos ¹ terbufos ¹	YES	NO	YES	NO	YES	NO
2B	Phenylpyrazoles (Fiproles)	fipronil ³	YES	NO	YES	NO	YES	NO
3A	Synthetic pyrethroids	bifenthrin ³	NO	YES	NO	YES	NO	YES
4A	Neonicotinoids	clothianidin ³ imidacloprid ³	NO	YES	NO	YES	NO	YES
4A/23	Neonicotinoid + tetramic acid	(imidacloprid + spirotetramat) ³	NO	YES	NO	YES	NO	YES
5	Spinosyns	spinetoram ² spinosad ²	NO	YES	NO	YES	NO	YES
28	Diamides	tetraniliprole ¹	NO	YES	NO	YES	NO	YES

*Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides

¹Products registered for banana weevil borer control

²Product registered for banana rust thrips control as bunch sprays only

³Products registered for banana weevil borer and banana rust thrips control

Guidelines

1. The resistance management strategy may start at any point in the product group rotation and planting may occur in any year of the strategy.
2. The product(s) used in any one year **should not be** followed by product(s) from the same Mode of Action (MOA) group in the following year.
3. Only products from the **YES** insecticide MOA groups shown in the diagram above **should be** applied for banana weevil borer control and/or banana rust thrips control in the same year. Where activity is specific to either banana weevil borer **or** banana rust thrips for an insecticide MOA group, then products from across these YES insecticide MOA groups may be applied in alternating years for their designated pest, but rotation of MOA groups between years for each pest should be maintained.
4. If products from **Group 1A or 1B** are being used for nematode control in a block of bananas, then products from these groups **should not be** used for banana weevil borer and/or banana rust thrips control in the following year.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



5. Where there is evidence of banana weevil borer and/or banana rust thrips resistance to a product or group of products, these **should not be** used again for banana weevil borer or banana rust thrips control until there has been use of products from other insecticide MOA groups for a period of at least 2 years.

STRATEGY B: Where products including controlled release (CR) formulations of imidacloprid are being used to control insects in bananas

MoA Group*	Chemical subgroup	Example chemical	Year 1 use	Year 2 use	Year 3 use	Year 4 use	Year 5 use	Year 6 use
1A or 1B	Carbamates Organophosphates	oxamyl ¹ or acephate ² cadusafos ¹ prothiofos ¹ terbufos ¹	NO	YES	NO	YES	NO	YES
2B	Phenylpyrazoles (Fiproles)	fipronil ³	YES	NO	YES	NO	YES	NO
3A	Synthetic pyrethroids	bifenthrin ³	NO	YES	NO	YES	NO	YES
4A	Neonicotinoids	CR imidacloprid ³	YES	YES	YES	NO	NO	NO
5	Spinosyns	spinetoram ² spinosad ²	YES	NO	YES	NO	YES	NO
28	Diamides	tetraniliprole ¹	NO	YES	NO	YES	NO	YES

8

*Refer: CropLife Australia Insecticide Resistance Management Review Group Mode of Action

Classification for Insecticides

¹Products registered for banana weevil borer control

²Product registered for rust thrips control as bunch sprays only

³Products registered for banana weevil borer and rust thrips control

Guidelines

1. The resistance management strategy may start at year 1 or year 4 in the product group rotation.
2. Controlled release imidacloprid provides 3 years control of banana weevil borer with one application at planting, so after the 3rd year, insecticide products from other Mode of Action (MOA) groups are to be used in rotation for at least 3 years for banana weevil borer and banana rust thrips control in a given block of bananas.
3. Alternative MOA groups are provided in these 3 years for control of banana rust thrips as soil or stem treatments or bunch sprays.
4. Only products from the **YES** Insecticide MOA Groups shown in the diagram above **should be** applied for banana weevil borer control and/or banana rust thrips control in the same year. Where activity is specific to either banana weevil borer **or** banana rust thrips for an insecticide MOA group, then

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



products from across these YES insecticide MOA groups may be applied in alternating years for their designated pest, but rotation of MOA groups between years for each pest should be maintained.

5. If products from Group 1A or 1B are being used for nematode control in a block of bananas, then products from these groups **should not be** used for banana weevil borer and/or banana rust thrips control in the following year.
6. Where there is evidence of banana weevil borer or banana rust thrips resistance to a product or group of products, these should not be used again for banana weevil borer and/or banana rust thrips control, until there has been use of products from other Insecticide Mode of Action groups for a period of at least 2 years.

Notes

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Brassica, brassica leafy vegetables
Insect(s)	Diamondback moth, Cabbage moth (<i>Plutella xylostella</i>)

Guidelines

1. To help prevent the development of resistance to any one specific active ingredient (see table below), observe the following instructions:
2. Use in accordance with the current IRMS for your region. For growers in the Lockyer Valley region, please refer to the [Lockyer Valley Diamondback Moth Insecticide Resistance Management Strategy](#):
 - a. Apply a specific active ingredient using a 'window' approach to avoid exposure of consecutive insect pest generations to the same mode of action. Multiple successive applications of a specific active ingredient are acceptable if they are used to treat a single pest generation.
 - b. Following a 'window' of a specific mode of action product, rotate to a 'window' of applications of effective insecticides with a different mode of action.
 - c. The total exposure period of any one mode of action 'active window' applied throughout the crop cycle (from seedling to harvest) should not exceed 50% of the crop cycle.
 - d. Incorporate IPM techniques into the overall pest management program.
 - e. Monitor insect populations for loss of field efficacy.
3. Always read and follow product labels and use the full recommended label rates of application. Some products place a limit on the number of times they can be applied per crop (see table below) and when they can be applied. 10
4. Monitor crops regularly and only apply insecticide when the pest threshold is reached.
5. When an insecticide with foliar activity on diamondback moth has been used as seed treatment or drench application in nursery production (as determined by label claims), rotate to an alternative mode of action insecticide for a period covering at least one generation of diamond back moth. This may require a minimum of 2 applications of alternate mode of action insecticides. Please refer to [CropLife's Nursery insecticide strategy](#) for further detail. ———
6. Ensure spray equipment is properly calibrated and achieves good coverage with appropriately sized spray droplets.
7. Time the application to the most susceptible life stage of the target pest.
8. To encourage beneficial insects, use *Bacillus thuringiensis* (Bt) sprays and avoid broad spectrum insecticides where possible, particularly early to mid-crop cycle.
9. Be cautious of using insecticide tank-mixes where both active ingredients control DBM as this strategy is generally not considered best practice for resistance management. Refer to the [IRAC International Insecticide Mixture Statement](#) for more information on this subject.
10. **DO NOT** re-treat a spray failure with a product from the same chemical group.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



11. Practice good crop hygiene to reduce DBM pressure - plant clean seedlings and incorporate crop residue as soon as practical after harvest

MoA Group*	Chemical subgroup	Active ingredient	Delivery method	No. applications permitted per crop per season
1A	Carbamates	methomyl (eg. Lannate® L), thiodicarb (eg. Larvin®)	Foliar	Not specified
2B	Phenylpyrazoles	fipronil (eg. Regent)	Foliar	4 per year within 8-week period
3A	Pyrethroids	synthetic pyrethroids (various – eg. Dominex® Duo, Karate® Zeon, Sumi-alpha® Flex, Trojan®)	Foliar	not specified
5	Spinosyns	spinetoram (Success® Neo), spinosad (eg. Entrust®)	Foliar	4
6	Avermectins	emamectin benzoate (eg. Proclaim® Opti)	Foliar	4 per any one crop
11A	<i>Bacillus thuringiensis</i>	Bacillus thuringiensis (eg. Dipel®, Xentari®)	Foliar	not specified
22A	Oxadiazines	indoxacarb (eg. Avatar® eVo)	Foliar	4
22A + 15	Oxadiazines + benzoylureas	indoxacarb + novaluron (Plemax®)	Foliar	3 (included as application of 22A)
23	Tetramic acid derivatives	spirotetramat (eg. Movento®)	Foliar	2 but 3 in brassica leafy vegetables
28	Diamides	chlorantraniliprole (eg. Coragen®), cyclaniliprole (Teppan®), flubendiamide (Belt®)	Foliar	3 and no more than 2 consecutive applications (including mixtures of chlorantraniliprole and thiamethoxam)
28 + 4A	Diamides + neonicotinoids	chlorantraniliprole + thiamethoxam (Durivo®)	Soil	1
30	Isoxazolines	broflanilide (Cimegra®), isocycloseram (Simodis®)	Foliar	2
UN	<i>Clitoria ternatea</i> extract	clitoria ternatea extract (Sero-X)	Foliar	not specified

11

* Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides

Notes

To ensure the most effective control of the pest:

- Product labels should at all times be carefully read and adhered to;
- Full recommended rates of registered insecticides should always be used; and

12. Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Canola, forage brassica
Insect(s)	Diamondback moth, Cabbage moth (<i>Plutella xylostella</i>)

Guidelines

For information, refer to the IPM guidelines: <https://ipmguidelinesforgrains.com.au/ipm-information/resistance-management-strategies/>.

Notes regarding the application of insecticide

1. To ensure the most effective control of the pest:
 - a. Product labels should at all times be carefully read and adhered to;
 - b. Full recommended rates of registered insecticides should always be used; and
 - c. Ensure good coverage of the target area to maximise contact.

To help prevent the development of resistance to any one specific active ingredient (see table below), observe the following instructions:

1. Apply a specific active ingredient using a 'window' approach to avoid exposure of consecutive insect pest generations to the same mode of action. Multiple successive applications of a specific active ingredient are acceptable if they are used to treat a single pest generation.
2. Following a 'window' of a specific mode of action product, rotate to a 'window' of applications of effective insecticides with a different mode of action.
3. The total exposure period of any one mode of action 'active window' applied throughout the crop cycle (from seedling to harvest) should not exceed 50% of the crop cycle.
4. Incorporate IPM techniques into the overall pest management program.
5. Consider the impact on beneficial insects. Consult the Cesar Australia Beneficials Chemical Toxicity table: <https://cesaraustralia.com/resources/beneficials-toxicity-table/>.
6. Monitor insect populations for loss of field efficacy.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



MoA Group*	Chemical subgroup	Active ingredient	No. applications permitted per crop per season	Crops
1A	Carbamates	methomyl (eg. Lannate® L), thiodicarb (eg. Larvin®)	not specified	Canola, rapeseed, brassicas
3A	Pyrethroids	synthetic pyrethroids (various – eg. Dominex® Duo, #Karate® Zeon, *Sumi-alpha® Flex, #Trojan®)	not specified	Canola, #forage brassicas, *fodder rape, *chou moellier
5	Spinosyns	spinetoram (Success® Neo)	2	Canola, forage brassicas
6	Avermectins	emamectin benzoate (eg. Affirm®)	2	Canola
11A	<i>Bacillus thuringiensis</i>	<i>Bacillus thuringiensis</i> (eg. Dipel®)	not specified	Oilseeds
28	Diamides	cyantraniliprole (Exirel®)	1	Canola
			2	Forage brassicas
31	Nucleopolyhedrovirus	NPV of H.zea or H.armigera – (eg. Gemstar®, Vivus® Max)	no limit but avoid season long use of low rates	Oilseeds
UN	Clitoria ternatea extract	clitoria ternatea extract (Sero-X)	not specified	Brassicas
UNM	Paraffinic spray oils	paraffinic oil (eg. Parachute®)	not specified when used in conjunction with Bt sprays	Canola

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)
Insect(s)

Cotton
All pests

Guidelines

For information refer to the current CottonInfo Cotton Pest Management Guide:

<https://www.cottoninfo.com.au/publications/cotton-pest-management-guide>

Notes

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Nursery e.g., vegetable seedlings, trees, ornamentals
Insect(s)	Various

Guidelines

1. Monitor regularly for insect activity. Where relevant consider the use of light, insect traps or other monitoring tools.
2. Only use insecticides when insect populations reach accepted threshold levels.
3. Always read and follow product labels and use the full recommended label rates of application. Some products place a limit on the number of times they can be applied per crop and when they can be applied.
4. Ensure the spray equipment is properly calibrated and achieves good coverage with appropriately sized spray droplets.
5. In the case of soil applied insecticides, apply once only. Apply using sufficient water that does not result in any drop point from the soil medium containing the seedling or plant. If watering is required between applications and planting, it should be done sparingly, only as required. **Avoid** water to run-through from the cells, bags etc.
6. When transplanting soil-treated seedlings or plants, ensure that the growing medium is fully transferred to the field with each seedling or plant.
7. **Rotate** between registered insecticides that have different modes of action.
8. Where possible **avoid applying consecutive applications** of insecticides that have the same mode of action within and between seasons or exceed the recommended maximum number of applications in a crop.
9. **Do not** follow a seed, seedling or soil treatment with a foliar application from the same mode of action group.
10. Time the foliar applications to the most susceptible life stage of the target pest.
11. **Do not** re-treat a spray failure with a product from the same mode of action group.
12. Avoid using insecticide tank-mixes where both active ingredients control the same insect pest as this strategy is generally not considered best practice for resistance management.
13. Practice good crop hygiene to reduce insect pressure e.g. removing severely infested seedling trays, plants or host weeds.
14. Nurseries supplying treated plants to commercial operations should clearly identify insecticides that have been applied and supply paperwork to the recipient accompanying the plant that indicates the rates and date. For treated seedlings the time of application should also be included, particularly if applied just prior to transplant in the field.

15

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Notes regarding the application of insecticides

1. Refer to the CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides.
2. There is known cross-resistance between some chemical groups e.g., **Groups 1A** and **1B**.
3. Seek advice from the manufacturers or government advisory services to determine local resistance levels for specific mode of action groups.
4. **Do not** exceed the maximum number of applications permitted on the insecticide label.
5. When using insecticides to control other pests consider the chemical group in relation to contributing to the resistance development of other insect pests.
6. When using insecticides to control insect pests consider the effect on beneficial insects and the potential to flare insect populations.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s) Pasture/Winter crops
Insect(s) Redlegged earth mite (RLEM) (*Halotydeus destructor*)

Guidelines

1. Determine the risk of RLEM for each paddock before making management decisions.¹
2. **Rotate** insecticide Groups.
3. **Do not** apply consecutive sprays of products from any one insecticide Group.
4. **Avoid** prophylactic pre-emergence/bare-earth sprays in autumn.

Crop stage	Group*	Chemical subgroup	Example chemical
Seed treatment (or in-furrow)	4A	neonicotinoids or	imidacloprid
	1B	organophosphates or	chlorpyrifos
	2B	phenylpyrazoles or	fipronil or isocycloseram (canola only)
	30	isoxazolines	
Bare Earth (Pre-emergent)	1B	organophosphates or	omethoate
	3A	synthetic pyrethroids	bifenthrin
Early post-emergence	1B	organophosphates or	chlorpyrifos
	3A	synthetic pyrethroids or	alpha-cypermethrin
	12A	diafenthiuron	diafenthiuron (Canola only)
Spring	1B	organophosphates or	omethoate
	3A	synthetic pyrethroids or	gamma-cyhalothrin
	12A	diafenthiuron	diafenthiuron (Canola only)

17

* Refer: *CropLife Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides*.

Further detailed information about registered products and chemical windowing strategies for RLEM is available at: https://ipmguidelinesforgrains.com.au/important/uploads/Science-behind-the-RLEM-IRMS-in-Australian-grains-and-pastures_October-2024.pdf

Monitoring and timing of sprays:

- Monitor RLEM activity carefully, particularly in the first 3-5 weeks after sowing.
- Only apply chemical if damage has reached economic threshold levels and/or the risk of RLEM is elevated.
- Use the hatch timing tool to estimate RLEM hatching in autumn.²
- Optimal timing of spring sprays can be calculated using the TIMERITE® tool.³ Follow the recommended best spray date and ensure RLEM is the dominant mite species.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Placement of sprays:

- Apply perimeter sprays where infestations are concentrated on the edge of paddocks.
- Spray the entire paddock when RLEM numbers exceed thresholds or risk is elevated. However, with pyrethroid products, recent research has shown that the risk of **Group 3A** resistance can be minimised through the application of foliar pyrethroids in a 50 m strip with 10 m spacing.⁴

Cultural practices:

- Decrease pasture foliage through grazing or cutting hay/silage to reduce RLEM numbers.
- Keep paddocks and fence-lines free of host weeds, such as capeweed.
- Sow crops and pastures that are more tolerant to RLEM, such as chickpeas.
- Encourage predator survival by judicious use of insecticides.
- Burn stubble prior to sowing to directly kill RLEM and its eggs.

Notes regarding the application of insecticides:

- There is known RLEM resistance to **Group 3A** and **Group 1B** insecticides in Western Australia, South Australia, and Victoria. Resistance is expected to expand across these regions.
- To ensure the most effective control of RLEM, follow the recommendations in the RLEM Best Management Practice Guide Revised November 2024
<https://ipmguidelinesforgrains.com.au/important/uploads/Redlegged-Earth-Mite-GRDC-24111322.pdf>
redlegged-earth-mite-grdc-20250130.pdf
 - Product labels should at all times be carefully read and adhered to;
 - Full recommended rates of registered insecticides should always be used; and
 - Ensure good coverage of the target area to maximise contact.

18

References

- ¹ The online RLEM seasonal risk tool developed by Cesar Australia considers multiple risks and management tactics
<https://cesaraustralia.shinyapps.io/rlemseasonalrisk/>
- ² The Hatch timing tool developed by Cesar Australia to aids early season monitoring and prediction of RLEM risk at crop establishment
<https://cesaraustralia.shinyapps.io/RLEM-hatch/>
- ³ TIMERITE® predicts the best spray window and ideal spray date in spring to control RLEM <https://www.wool.com/timerite/>
- ⁴ [Strip spraying delays pyrethroid resistance in the redlegged earth mite, Halotydeus destructor: a novel refuge strategy - Maino - 2021 - Pest Management Science - Wiley Online Library](#)

Crop(s)

Pome fruit

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Insect(s) **Two-spotted mite (*Tetranychus urticae*), European red mite (*Panonychus ulmi*)**

Guidelines

1. Best management practice is to make **no more than one application** from each registered miticide group per season. Rotate registered miticides that have different mode of action (i.e. **Group 6, Group 10A, Group 10B, Group 12B, Group 12C, Group 13, Group 20B, Group 20D, Group 21A, Group 23, and Group 25A**).
2. For miticides that have the same mode of action (e.g. **Group 21A**) do not use sequential applications between seasons. Sequential applications include from the end of one season to the beginning of the next.

Group*	Chemical subgroup	Example chemical
6	Avermectins, milbemycins	abamectin, milbemectin
10A	Clofentezine, hexythiazox	clofentezine, hexythiazox
10B	Etoxazole	etoxazole
12B	Organotin miticides	fenbutatin oxide
12C	Propargite	propargite
20B	Acequinocyl	acequinocyl
20D	Bifenazate	bifenazate
21A	METI acaricides	fenpyroximate, tebufenpyrad
23	Tetronic and Tetramic acid derivatives	spiromesifen
25A	Beta-ketonitrile derivatives	cyflumetofen

* Refer: *CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides*.

19

Notes

- Miticides should be used as part of an Integrated Mite Control (IMC) program.
- Mite levels should be monitored, and thresholds utilised before deciding to make miticide applications.
- Where practicable, predatory mites should be incorporated into an IMC program.
- When using insecticides or miticides to control other pests of pome fruit, such as codling moth, lightbrown apple moth and woolly aphid, consider the chemical group and the potential impact it may have on resistance development of mite pests.
- When using insecticides or miticides to control other pests of pome fruit, consider the effect on beneficial insects and the potential to flare mite populations.
- For more information refer to the current [NSW Orchard Plant Protection Guide](#).

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s) **Potato**
Insect(s) **Potato tuber moth / Tomato leafminer (*Phthorimaea operculella*)**

Guidelines

1. Monitor pest levels and **do not** spray unless pest thresholds are exceeded.
2. **Rotate** insecticide groups and **do not** apply more than two consecutive applications of products with the same Mode of Action.
3. Integrate both chemical and non-chemical means of control as part of the overall control strategy. Examples are the use of predators/parasites and relevant cultural practices (crop hygiene, rotation of planted areas, and strategic time of planting).

Group*	Chemical subgroup	Example chemical
1A	Carbamates	carbaryl, methomyl
1B	Organophosphates	acephate, ,
3A	Pyrethroids	permethrin
5	Spinosyns	spinosad, spinetoram
28	Diamides	chlorantraniliprole, flubendiamide, cyantraniliprole

* Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides.

21

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Always read and follow product labels and use the full recommended label rates of application; and
- b) Ensure the spray equipment is properly calibrated and achieves good coverage of target area with appropriately sized spray droplets.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Sorghum, maize, summer & winter grain legumes
Insect(s)	Heliothis/Cotton bollworm/Native budworm (<i>Helicoverpa</i> spp.)

Guidelines

- To help prevent the development of resistance to any specific active ingredient (see table below), observe the following instructions:
 - Use in accordance with the current IRMS for your region.
 - Apply a specific active ingredient using a “window” approach to avoid exposure of consecutive insect pest generations to the same mode of action. Multiple successive applications of a specific active ingredient are acceptable if they are used to treat a single pest generation.
 - Following a ‘window’ of a specific mode of action product, rotate to a ‘window’ of applications of effective insecticides with a different mode of action.
 - The total exposure period of any one mode of action ‘active window’ applied throughout the crop cycle (from seedling to harvest) should not exceed 50% of the crop cycle.
 - Incorporate IPM techniques into the overall pest management program.
 - Monitor insect populations for loss of field efficacy.
- Always read and follow product labels. Some products place a limit on the number of times they can be applied per crop (see table below) and when they can be applied. 22
- Monitor crops regularly and only apply insecticide when the pest threshold is reached.
- Ensure spray equipment is properly calibrated and achieving good coverage with appropriately sized spray droplets.
- Time the application to the most susceptible life stage of the target pest.
- To encourage beneficial insects, use *Bacillus thuringiensis* (Bt) or NPV sprays and avoid broad spectrum insecticides where possible, particularly early to mid-crop cycle.
- Be cautious of using insecticide tank-mixes where both active ingredients control *Helicoverpa* spp. as this strategy is generally not considered best practice for resistance management. Refer to document [IRAC International Insecticide Mixture Statement](#) for more information on this subject.
- Do not** re-treat a spray failure with a product from the same chemical group.
- Practice effective pupae busting as soon as practicable after harvest.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



10. The Modes of Action (Groups) and registered insecticides for control of Heliothis/Cotton bollworm/Native budworm (*Helicoverpa* spp.) are listed below:

MoA Group*	Chemical subgroup	Active ingredient	No. applications permitted per crop per season	Crops
1A	Carbamates	methomyl (eg. Lannate® L), thiodicarb (eg. Larvin®)	not specified	All cereal grains, oilseed, pulses
3A	Pyrethroids	synthetic pyrethroids (various – eg. Dominex® Duo, Karate® Zeon, Sumi-alpha® Flex, Trojan®)	not specified	All cereal grains, oilseed, pulses
5	Spinosyns	spinetoram (Success® Neo)	2	All pulses
5 + 18	Spinosyns + diacylhydrazines	spinetoram + methoxyfenozide (Intrepid Edge®)	Chickpeas: 1	Chickpeas
			Mung beans: 2	Mung beans
6	Avermectins	emamectin benzoate (eg. Affirm®)	2	All pulses
6 + 4A	Avermectins + neonicotinoids	emamectin benzoate + acetamiprid (Skope®)	2	Summer & winter pulses except field peas and lupins
11A	<i>Bacillus thuringiensis</i>	<i>Bacillus thuringiensis</i> (eg. Dipel®)	not specified	All cereal grains, oilseed, pulses
22A	Oxadiazines	indoxacarb (eg. Steward® EC)	1	chickpea, faba bean, mung bean, soybean, azuki bean
28	Diamides	chlorantraniliprole (eg. Vantacor®)	2	All pulses
31	Nucleopolyhedrovirus	NPV of H.zea or H.armigera – (eg. Gemstar®, Vivus® Max)	no limit but avoid season long use of low rates	All cereal grains, oilseed, pulses
UNM	Paraffinic spray oils	paraffinic oil (eg. Parachute®)	2 alone or in combination with NPV's	All pulses, oilseeds

* Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides.

¥ Refer: Registered product label

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Notes

- For more information refer to the IPM Guidelines *H. armigera* RMS for Australian grains:
<https://ipmguidelinesforgrains.com.au/ipm-information/resistance-management-strategies/#heli>
- Consider the impact on beneficial insects. Consult the Cesar Australia beneficial insects table:
<https://cesaraustralia.com/resources/beneficials-toxicity-table/>
- Queensland DPI resistance monitoring updates: <https://thebeatsheet.com.au/key-pests/helicoverpa-management/>

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Strawberries / Ornamentals
Insect(s)	Two-spotted mite (<i>Tetranychus urticae</i>)

Guidelines

1. Monitor mite activity and treat infestations before thresholds are reached, i.e. spray earlier rather than later. Seek advice on local threshold levels.
2. **Do not** apply sequential applications of products from any one chemical group. This does not apply to various oils or sulphur as they are considered to have a lower resistance risk.
3. Preferably products with the same Mode of Action should not be used more than twice in a growing season. This does not apply to various oils or sulphur as they are considered to have a lower resistance risk.
4. Incorporate the use of predatory mites for the control of this pest wherever possible.

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
5. Full recommended rates of registered insecticides should always be used; and
 6. Ensure good coverage of the target area to maximise contact.

Continued on next page.

25

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



MoA Group*	Chemical subgroup	Active ingredient	No. applications permitted per crop per season	Crops
1B	Organophosphates	Dimethoate	not specified	Ornamentals (refer labels)
3A	Pyrethroids Pyrethrins	Bifenthrin	2	ornamentals
6	Avermectins Milbemycins	Abamectin, Milbemectin	2	strawberries, ornamentals
10A	Clofentezine Hexythiazox Pyridaben	Clofentezine Hexythiazox Pyridaben	1	Strawberries, Ornamentals
12B	Organotin miticides	Fenbutatin Oxide	not specified	strawberries, ornamentals
12C	Propargite	Propargite	A maximum of 3 sprays applied over a period of 3 to 4 weeks	strawberries, ornamentals
20D	Bifenazate	Bifenazate	2	strawberries
25A	Beta-ketonitrile derivatives	Cyflumetofen	2	strawberries, ornamentals
UN	Sulphur	Sulphur (S) as Wetttable Sulphur	not specified	ornamentals
UNM	Emulsifiable Botanical Oils, Paraffinic Oils	Emulsifiable Botanical Oils, Orange oil, Paraffinic Oils	no more than three in an 8 week period	strawberries, ornamentals

26

Expert Committee on Insecticide Resistance guidance recommend a maximum of 2 non-sequential applications per season. See Guidelines, Point 2

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Sweet corn
Insect(s)	Corn earworm (<i>Helicoverpa armigera</i>) aka Heliothis

Guidelines

1. If Fall armyworm is the dominant pest, refer to the Fall armyworm (*Spodoptera frugiperda*) strategy. Many insecticides used for Heliothis control will also place a selection pressure on Fall armyworm.
2. The critical stage of infestation is during silking. Even low levels of Heliothis infestation are unacceptable at the silking stage. As sweet corn is less attractive to Heliothis before flowering and it is picked soon after silking is completed, there is a relatively short period of protection required.
3. Control of Heliothis at the tasselling stage (occurs prior to silking stage) can be important in some regions as the tassel can act as a nursery for Heliothis, which can then move onto the young developing cobs. Control of Heliothis at this stage is not as difficult as at the silking stage. Use of biological insecticides, Bt and Nuclear Polyhedrosis Virus (NPV), in the early stages of crop development is encouraged.
4. Monitor crops regularly, at least weekly during silking and **do not** spray unless pest thresholds are exceeded.
5. Labels of new products place a limit on the number of applications. If further control is required on one planting, chemicals from different mode of action groups within the same window should be used.
6. **Do not** retreat a spray failure with a product from the same chemical group.
7. **Do not** use mixtures of insecticides for controlling Heliothis.
8. Cultivation after harvest to destroy pupae will greatly assist in managing Heliothis.
9. Seek local advice on pest incidence and on the risk of resistance developing from insecticide programs used to control Heliothis in crops other than sweet corn.
10. To help prevent the development of resistance to any one specific active ingredient (see table below), observe the following instructions:
 - a. Use in accordance with the current IRMS for your region;
 - b. Apply a specific active ingredient using a “window” approach to avoid exposure of consecutive insect pest generations to the same mode of action. Multiple successive applications of a specific active ingredient are acceptable if they are used to treat a single insect generation;
 - c. Following a “window” of a specific mode of action product, rotate to a “window” of applications of effective insecticides with a different mode of action;

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



- d. The total exposure period of any one mode of action “active window” applied throughout the crop cycle (from seedling to harvest) should not exceed 50% of the crop cycle;
- e. Incorporate IPM techniques into the overall pest management program; and
- f. Monitor insect populations for loss of field efficacy.

MoA Group*	Chemical subgroup	Active ingredient	Delivery method	No. applications permitted per crop per season
1A	Carbamates	methomyl (eg. Lannate® L), thiodicarb (eg. Larvin®)	Foliar	not specified
3A	Pyrethroids	synthetic pyrethroids (various – eg. Dominex® Duo, Sumi-alpha® Flex)	Foliar	not specified
5	Spinosyns	spinetoram (Success® Neo), spinosad (eg. Entrust®)	Foliar	4
6	Avermectins	emamectin benzoate (eg. Proclaim® Opti)	Foliar	4
11A	<i>Bacillus thuringiensis</i>	<i>B. thuringiensis subsp. kurstaki</i> (eg. Dipel®)	Foliar	not specified
18	Diacylhydrazines	Methoxyfenozide (Permit No. 84531, 30 June 2030)	Foliar	3
22A	Oxadiazines	indoxacarb (eg. Avatar® eVo)	Foliar	3
28	Diamides	chlorantraniliprole (eg. Coragen®),	Foliar	3 and no more than 2 consecutive applications
31	Nucleopolyhedrovirus	NPV of <i>H. zea</i> or <i>H. armigera</i> (eg. Gemstar®, Vivus® Max)	Foliar	not specified

28

* Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Region	January	February	March	April	May	June	July	August	September	October	November	December																
North Queensland	No Crop				Vegetative phase		Emamectin Benzoate		Methomyl, Thiodicarb, SP's			Chlorantraniliprole			Spinetoram			No Crop										
Heliiothis pressure	L	L	L	L	M	H	H	H	H	M	L	L	L	L	M	H	H	H	H	H	H	M	L	L				
South East Queensland	Spinetoram		Chlorantraniliprole				Methomyl, Thiodicarb, SP's			No Crop						Vegetative phase			Emamectin Benzoate				Spinetoram					
Heliiothis pressure	H	H	H	H	H	H	M	M	L	L	L	L	L	L	L	L	L	M	M	H	H	H	H	H				
Central NSW / Northern Victoria	Spinetoram				Chlorantraniliprole				Methomyl Thiodicarb SP's		No crop							Vegetative phase			Emamectin Benzoate							
Heliiothis pressure	H	H	H	H	H	M	M	L	L	L	L	L	L	L	L	L	L	L	M	M	M	M	H	H				
Tasmania	Chlorantraniliprole			Spinetoram			No crop																		Emamectin Benzoate			
Heliiothis pressure	M	M	M	M	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L				
In all regions	Nuclear Polyhedrosis Viruses (NPV's), <i>Bacillus thuringiensis</i> (Bt) and Methomyl at the ovicidal rate can be used season long with no resistance management implications																											

H: High pressure period; M: medium pressure period; L: Low pressure period

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Tomato
Insect(s)	Heliothis / Tomato budworm (<i>Helicoverpa</i> spp.)

Guidelines

1. Monitor pest levels and **do not** spray unless pest thresholds are reached.
2. Use an integrated pest control approach where both chemical and non-chemical measures are adopted as part of the overall strategy. Examples are the use of predators/parasites and relevant cultural practices (crop hygiene, rotation of planted areas, and strategic time of planting).
3. Seek local advice on pest incidence and the risk of resistance development from insecticide programs used to control *Heliothis* in other crops or to control other pests.
4. When using insecticides/miticides to control other pests on tomato, consider the chemical group in relation to contributing to resistance development of *Heliothis*.
5. Avoid using insecticides from the same chemical group against consecutive generations of *Helicoverpa* spp. or other pests, as this will increase the selection pressure. This does not apply to *Bacillus thuringiensis* (Bt) or Nucleopolyhedroviruses (NPV) products, as they are considered to have a lower resistance risk. To encourage beneficial insects, use *Bacillus thuringiensis* (Bt) or Nucleopolyhedroviruses (NPV) sprays and avoid broad spectrum insecticides where possible, particularly early to mid-crop cycle.
6. Do not re-spray a crop in the same season where a failure (which is known or unknown) has occurred using the same insecticide or other active ingredient from the same chemical group.
7. The Modes of Action (Groups) and registered insecticides for control of *Heliothis*/Tomato budworm (*Helicoverpa* spp.) are listed below:

(Continued on next page)

30

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



MoA Group*	Chemical subgroup	Active ingredient	Delivery method	No. applications permitted per crop per season
1A	Carbamates	methomyl (eg. Lannate® L), thiodicarb (eg. Larvin®)	Foliar	not specified
3A	Pyrethroids	synthetic pyrethroids (various – eg. Dominex® Duo, Karate® Zeon, Sumi-alpha® Flex, Trojan®)	Foliar	not specified
5	Spinosyns	spinetoram (Success® Neo), spinosad (eg. Entrust®)	Foliar	4
6	Avermectins	emamectin benzoate (eg. Proclaim® Opti)	Foliar	4
11A	<i>Bacillus thuringiensis</i>	<i>B.thuringiensis subsp. kurstaki</i> (eg. Dipel®)	Foliar	not specified
18	Diacylhydrazines	methoxyfenozide (eg. Prodigy®)	Foliar	not specified
22A	Oxadiazines	indoxacarb (eg. Avatar® eVo)	Foliar	3 and no more than 2 consecutive applications
22A + 15	Oxadiazines + benzoylureas	indoxacarb + novaluron (Plemax®)	Foliar	3 and no more than 2 consecutive applications
28	Diamides	chlorantraniliprole (eg. Coragen®), flubendiamide (eg. Belt®)	Foliar	3 and no more than 2 consecutive applications (including mixtures of chlorantraniliprole and thiamethoxam)
28	Diamides	cyantraniliprole (Benevia®)	Foliar	2
28 + 12A	Diamides + diafenthiuron	cyantraniliprole + diafenthiuron (Minecto Forte®)	Foliar	2
28 + 4A	Diamides + neonicotinoids	chlorantraniliprole + thiamethoxam (Durivo®)	Soil	1
31	Nucleopolyhedrovirus	NPV of H.zea or H.armigera (eg. Gemstar®, Vivus® Max)	Foliar	not specified
UN	<i>Clitoria ternatea</i> extract	clitoria ternatea extract (Sero-X)	Foliar	not specified

31

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s) **Turf**
Insect(s) / Mite(s) **Various**

Guidelines

The Modes of Action (groups) and registered miticide and insecticides in turf are listed below.

Miticides

Group*	Chemical subgroup	Example chemical	Product type
3A	Pyrethroids	beta-cyfluthrin bifenthrin	Miticide/Insecticide
6	Avermectins	abamectin	Miticide
10A	Clofentezine	clofentezine	Miticide
10B	Etoxazole	etoxazole	Miticide
12A	Diafenthiuron	diafenthiuron	Miticide
UN	Azadirachtin	azadirachtin	Miticide

Insecticides

Group*	Chemical subgroup	Example chemical	Product type
1A	Carbamates	bendiocarb	Insecticide
1B	Organophosphates	diazinon maldison (malathion) trichlorfon	Insecticide
4A	Neonicotinoids	clothianidin imidacloprid thiamethoxam	Insecticide
2B	Phenylpyrazoles (Fiproles)	fipronil	Insecticide
15	Benzoylureas	novaluron	Insecticide
7C	Pyriproxyfen	pyriproxyfen	Insecticide
11A	<i>Bacillus thuringiensis</i> and the insecticidal proteins they produce	<i>Bacillus thuringiensis</i>	Insecticide
20A	Hydramethylnon	hydramethylnon	Insecticide
22A	Oxadiazines	indoxacarb	Insecticide
22B	Semicarbazones	metaflumizone	Insecticide
28	Diamides	chlorantraniliprole cyantraniliprole tetraniliprole	Insecticide
UN	Azadirachtin	azadirachtin	Insecticide

* Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Guidelines

1. Monitor regularly for insect/mite activity. Where relevant consider the use of light, insect traps or other monitoring tools, including flushing with pyrethrum, soap, or salt solutions.
 2. Use insecticides when insect populations reach accepted threshold levels.
 3. For certain insect pests, preventative applications may be required to avoid the buildup of insect populations.
 4. Always read and follow product labels and use the full recommended label rates of application. Some products place a limit on the number of times they can be applied per season/year and when they can be applied.
 5. Ensure the spray equipment is properly calibrated and achieving good coverage with appropriately sized spray droplets.
 6. **Do not** apply if heavy rains or storms that are likely to cause runoff are forecast. Read individual labels for additional advice.
1. **Rotate** between registered insecticides that have different modes of action.
 2. Where possible **avoid applying consecutive applications** of insecticides that have the same mode of action within and between seasons or exceed the recommended maximum number of applications in a season/year.
 3. Time the applications to the most susceptible life stage of the target pest.
 4. **Do not** re-treat a spray failure with a product from the same mode of action group.
 5. Avoid using insecticide tank-mixes where both active ingredients control the same insect pests as this strategy is generally not considered best-practice for resistance management.

34

Mite – additional information

1. Monitor mite activity and treat infestations as soon as detected i.e. spray earlier rather than later. Seek advice on local threshold levels.
2. Avoid sequential applications of products from any one mode of action group.
3. Preferably products with the same Mode of Action should not be used more than twice during the main growing season.

Notes

1. Refer to The CropLife Australia Expert committee on Insecticide Resistance Mode of Action Classification for Insecticides.
2. There is known cross-resistance between some chemical groups e.g. **Groups 1A and 1B**.
3. Seek advice from the manufacturers and/or government advisory services to determine local resistance levels for specific mode of action **Groups**.
4. **Do not** exceed the maximum number of applications permitted on the insecticide/miticide label.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



5. When using insecticides to control other pests, consider the chemical group in relation to contributing to resistance development of other insect/mite pests.
6. When using insecticides to control pests, consider the effect on beneficial insects and the potential to flare insect/mite populations.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Various
Insect(s)	Cotton/Melon aphid (<i>Aphis gossypii</i>), Green peach aphid (<i>Myzus persicae</i>)

Guidelines

1. **Rotate** between registered insecticides that have different modes of action (e.g., **Groups 1, 4, 9, 12A** (cotton crop only), **23, 28** and **29**).
2. **DO NOT** apply consecutive applications of **Group 4C** insecticides for control of aphids.
3. **Do not** apply more than two sequential applications of insecticides that have the same mode of action within and between seasons or exceed the recommended maximum number of applications in a crop.
4. With post-spray monitoring, **do not** re-treat a spray failure with a product from the same mode of action group.
5. **Do not** follow a seed/seedling/soil treatment with a foliar application from the same mode of action group.
6. The Modes of Action (Groups) and registered insecticides for control of cotton/melon aphid and/or green peach aphid are listed below:

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Group*	Chemical subgroup	Example active ingredient
1A	Carbamates	pirimicarb
1B	Organophosphates	dimethoate
3A	Pyrethroids and Pyrethrins	tau-fluvalinate
4A	Neonicotinoid	imidacloprid
4C*	Sulfoximines	sulfoxaflor
8A	Alkyl halides	ethyl formate
9B	Pyridine azomethine derivatives	pymetrozine
9D	Pyropenes	afidopyropen
12A	Diafenthion	diafenthion
23	Tetronic and Tetramic acid derivatives	spirotetramat
28	Diamides	cyantraniliprole
29	Flonicamid	flonicamid
36	Pyridazine pyrazolecarboxamides	dinpropylpyridaz
UN	Sulphur	sulphur
UNF		<i>Beauveria bassiana</i>
UNM		Paraffinic oil

* Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Notes

- There is known cross-resistance between **Groups 1A and 1B**. Rotate between **Group 1** and **Group 4, 9B, 9D, 12A, 23, 28 and 29**.
- Consecutive applications of a **Group 4A** product may be made only if no other effective option is available – either because:
 - no other group is registered in the crop; or
 - the target pest is resistant to the other Groups.
- Seek advice from the manufacturers and/or government advisory services to determine local resistance levels for particular mode of action Groups.
- **Do not** exceed the maximum number of applications permitted on the insecticide label.
- When using insecticides/miticides to control other pests, consider the chemical group in relation to contributing to resistance development of Cotton/Melon Aphid and Green Peach Aphid.
- When using insecticides/aphicides to control other pests consider the effect on beneficial insects and the potential to flare aphid populations.
- For more information about insecticide resistance in the green peach aphid refer to <http://cesaraustralia.com/latest-news/all/insecticide-resistance-in-the-green-peach-aphid>.
- For more detail on resistance management of aphids in cotton refer to the current [Cotton Pest Management Guide](#) or for more detail on resistance management for Green Peach Aphid in grain refer to <https://ipmguidelinesforgrains.com.au/ipm-information/resistance-management-strategies/>, <https://grdc.com.au/resources-and-publications/all-publications/publications/2024/green-peach-aphid-best-management-practice-guide>, and <https://thebeatsheet.com.au/key-pests/aphids>.

38

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Various
Insect(s)	Fall armyworm (<i>Spodoptera frugiperda</i>)

Guidelines

List of active constituents approved for use under permits or registered by the Australian Pesticides and Veterinary Medicines Authority (APVMA) as of March 2026:** Refer to the APVMA's Agricultural And Veterinary Permits Search website (<https://portal.apvma.gov.au/permits>) to ensure permit is still active.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



MoA Group	Chemical sub-group	Active ingredient	No. applications permitted per crop per season	Permit number*	Crops
5	Spinosyns	spinetoram	3	PER93550 to 31.12.26	Ginger
5	Spinosyns	spinosad	4	PER89870 to 31.10.30	Various vegetables, berryfruit, coffee, tropical and subtropical fruit (inedible peel), ornamentals, sweet corn
5	Spinosyns	spinetoram	4	PER89241 to 31.1.28	capsicum, sweet corn
			2	PER93482 to 30.9.26	Maize, cereals, sorghum (grain), millet
5 + 18	Spinosyns + diacylhydrazines	spinetoram + methoxyfenozide	2	PER95222 to 31.10.28	Maize, popcorn
6	Avermectins	emamectin benzoate	4	PER89263 to 31.1.28	Capsicum
			4	PER95680 to 31.1.28	Ginger
22A	Oxadiazines	indoxacarb	3	PER93815 to 31.12.28	Sweet corn
			2	PER93488 to 30.9.26	Maize cereals
			1	PER90761 to 28.2.27	Linseed
28	Diamides	chlorantraniliprole	3	PER89259 to 31.1.28	Capsicum, sweet corn
			2	PER96040 to 30.4.27	Ginger

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



			3	PER95049 to 3.1.27	Cane berries
			2	PER86014 to 31.7.26	Peanuts
31	Nucleopolyhedrovirus	SfMNPV	5	PER91477 to 31.3.27	Cereal grains, oilseed, pulses, fodder and forage crops
			10		Cotton, sweet corn, corn, root and tuber vegetables, legume vegetables, ornamentals
			5	PER90820 to 31.3.27	Cereal grains, oilseed, pulses, fodder and forage crops
			10		Cotton, sweet corn, root and tuber vegetables, legume vegetables, fruiting vegetables (other than cucurbits), Leafy vegetables (inc. brassica leafy vegetables), ornamentals
Various			Various	PER91928 to 30.9.30 PER91923 to 31.5.27 PER91806 to 30.11.26	Nursery stock (non-food)

**Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides*

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Guidelines

1. An Integrated Pest Management (IPM) approach should be adopted in the production system to help manage this pest, with focus on cultural methods and the preservation of beneficial arthropods (insects and spiders):
 - a. This includes regular crop monitoring (at least 2 times per week) to determine incidence x crop damage and the impact of beneficial arthropods.
 - b. Consideration should also be given to the impact of prevailing weather conditions on the rate of pest development in the field.
2. **Avoid** sequential plantings of preferred crops, e.g. corn, sorghum, sugarcane, as this will increase local populations of fall armyworm.
3. Management of crop residues/volunteer plants before planting and after harvest also helps reduce local populations of fall armyworm.
4. Where possible, **avoid** the use of broad spectrum foliar applied insecticides in the production system for both larvae and moth control. If broad-spectrum insecticides are to be used, apply at timings when preservation of beneficial species is less likely to be important – i.e. at end of growing season.
5. Consider controlling moths using light or attractant traps or sprays and encourage micro-bat habitat (natural or artificial roosting sites) adjacent to production areas.
6. In situations where insecticides are required, consider beneficial arthropods when making spray decisions.
7. When applying insecticides to this pest, key considerations should be given to:
 - a. Apply insecticides only when needed based on economic thresholds;
 - b. **Target early instar stages** (hatching larvae) of the pest before they become entrenched in the crop e.g. lower whorl of maize, sweet corn or grain sorghum;
 - c. Use a **medium spray quality** to ensure sufficient droplets cover the spray target to ensure the larvae ingest a lethal dose of insecticide; and
 - d. Use a well calibrated, functioning boom spray with appropriate water rate for the target crop to ensure optimum spray coverage.
8. Use the recommended insecticide rates as stipulated on the relevant APVMA Emergency Use Permit or label.
9. Use a recommended adjuvant if stipulated on the relevant APVMA Emergency Use Permit or label.
10. Inspect the performance of the insecticide 3-4 days after application. Always document the effectiveness of each insecticide application and never re-spray a failure with an insecticide with the same mode of action. Inform your local reseller or agronomist of any spray failures. Internationally, known resistance has occurred to the following MoA groups: Carbamates (Group 1A); Organophosphates (Group 1B); Pyrethroids (Group 3); Bacillus thuringiensis and Cry1F protein (Group 11A).

42

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



11. When using selected insecticides in-crop targeting fall armyworm, the following resistance management strategy guidelines should be implemented:
 - a. Do not treat successive generations with products of the same MOA;
 - b. The total exposure period of any one MOA insecticide applied throughout the crop cycle (from seedling to harvest) **should not exceed 50% of the crop cycle**;
 - c. Abide by the individual label recommendation for maximum **number of allowable applications per crop per season**;
 - d. Abide by individual label recommendation for the **minimum reapplication interval** and always use the **full recommended label rates**;
 - e. Where possible, an Area Wide Management strategy should be adopted where the same MOA insecticides are used **by all growers** in the **same time period**; and
 - f. As the industry learns more about how to manage this pest, this Strategy may be updated and regional-specific strategies may be developed. Check the CropLife Resistance Management [website](#) to ensure you are following the most up to date fall armyworm strategy.
12. Useful fall armyworm reference documents are available:
 - <https://portal.apvma.gov.au/permits>: search for APVMA insecticide permits to use on FAW.
 - <https://www.cottoninfo.com.au/insect-and-mite-management>
 - <https://thebeatsheet.com.au/fall-armyworm-should-you-be-concerned/>
 - <https://irac-online.org/pests/spodoptera-frugiperda/>
 - <https://irac-online.org/new-guidelines-on-ipm-irm-for-fall-armyworm-in-s-african-maize/>
 - <https://grdc.com.au/resources-and-publications/resources/fall-armyworm>

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Various
Insect(s)	Mites

Guidelines

13. For Pome fruits, Strawberries/ornamentals and Redlegged earth mite see specific CropLife strategies.
1. Monitor mite activity, accurately identify species, and treat infestations before thresholds are reached, i.e. spray earlier rather than later. Seek advice on local threshold levels.
2. **Do not** apply sequential applications of products from any one chemical group.
3. Preferably products with the same Mode of Action should not be used more than twice in a growing season.
4. Incorporate the use of predatory mites and broader IMP techniques for the control of this pest wherever possible.
5. Incorporate IPM techniques into the overall pest management program. For more information see <https://www.horticulture.com.au/growers/help-your-business-grow/research-reports-publications-fact-sheets-and-more/vg16067/>

Notes regarding the application of miticides:

1. Refer to The CropLife Australia Expert committee on Insecticide Resistance Mode of Action Classification for Insecticides.
2. There is known cross-resistance between some chemical groups e.g. **Groups 1A and 1B**.
3. Seek advice from the manufacturers and/or government advisory services to determine local resistance levels for specific mode of action Groups.
4. Do not exceed the maximum number of applications permitted on the insecticide/miticide label.
5. When using insecticides to control other pests, consider the chemical group in relation to contributing to resistance development of other insect/mite pests.
6. When using insecticides to control pests, consider the effect on beneficial insects and the potential to flare insect/mite populations
7. To ensure the most effective control of the pest:
 - a. Product labels should at all times be carefully read and adhered to;
 - b. Full recommended rates of registered insecticides should always be used; and
 - c. Ensure good coverage of the target area to maximise contact.

44

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s) Insect(s)	Various Silverleaf whitefly (<i>Bemisia tabaci</i>)
----------------------	--

Guidelines

1. Monitor pest numbers and apply control measures before adult populations reach high levels.
2. Select registered insecticide control measures according to the primary growth stage of the pest, the infestation level and the age and type of crop.
3. In cotton, spray decisions should be based on the Silverleaf Whitefly threshold matrix. Refer to the current [Cotton Pest Management Guide](#) for further details.
4. Where possible, utilise selective insecticides during the early stages of crop development to minimise the impact on beneficial insects.
5. Rotate between registered insecticides that have different modes of action (see table below).
6. **Do not** apply more than two consecutive applications of insecticides that have the same Mode of Action within and between seasons.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



7. The Modes of Action (groups) and registered insecticides for control of Silverleaf Whitefly are listed below:

Group*	Chemical subgroup	Active ingredient
1B	Organophosphate	acephate
3A	Pyrethroids	bifenthrin
4A	Neonicotinoids	acetamiprid, clothianidin, dinotefuran, imidacloprid, thiamethoxam
4D	Butenolides	flupyradifurone
7C	Pyriproxyfen	pyriproxyfen
9D	Pyropenes	afidopyropen
12A	Diafenthiuron	diafenthiuron
16	Buprofezin	buprofezin
23	Tetronic and Tetramic acid derivatives	spirotetramat
28	Diamides	cyantraniliprole
28 + 4A	Diamides + Neonicotinoids	cyantraniliprole + thiamethoxam
29	Flonicamid	flonicamid
36	Pyridazine pyrazolecarboxamides	dimpropyridaz
UN		clitoria ternatea extract
UNF	<i>Beauveria bassiana</i> strains	<i>Beauveria bassiana</i> strain PPRI 5339
UNM	Mineral oil	paraffinic oil

*Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides

8. Seek advice from the manufacturers and/or government advisory services to determine local resistance levels for particular mode of action Groups.
9. **Do not** exceed the maximum number of applications permitted on the insecticide label.
10. When using insecticides to control other pests, consider the chemical group in relation to contributing to resistance development of Silverleaf Whitefly.
11. When using insecticides to control other pests consider the effect on beneficial insects and the potential to flare Silverleaf Whitefly populations.

Notes

Not all chemical groups listed have registered products available in all crops affected by Silverleaf Whitefly. Only use products registered for use in crop to be treated.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Cultural practices

1. In vegetable crops, ensure seedlings are free of pests prior to transplanting. Inspect transplants carefully upon arrival for whitefly eggs, nymphs and adults.
2. Control alternate weed hosts of Silverleaf Whitefly 2-3 weeks before planting to reduce early population levels.
3. Clean-up crop residues:
 - a. Where moderate population levels remain after harvest, apply a registered insecticide or oil treatment effective against adults.
 - b. Plough in crops within 2-3 days of application to kill all remaining nymphs on crop foliage to reduce pest migration into new plantings.

Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Product labels should at all times be carefully read and adhered to;
- b) Full recommended rates of registered insecticides should always be used; and
- c) Ensure good coverage of the target area to maximise contact.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Various
Insect(s)	Western flower thrips (<i>Frankliniella occidentalis</i>)

Guidelines

1. Monitor regularly for **Western flower thrips** (WFT) activity. WFT can cause damage to crops by feeding on them and can spread **tomato spotted wilt virus (TSWV)** in crops.
2. Only use insecticides when WFT populations reach accepted threshold levels.
3. Always read and follow product labels and use the full recommended label rates of application. Some products place a limit on the number of times they can be applied per crop and when they can be applied. DO NOT exceed the recommended maximum number of applications in a crop.
4. Ensure the spray equipment is properly calibrated and achieves good coverage with appropriately sized spray droplets.
5. Rotate between registered insecticides that have different modes of action.
6. Do not follow a seedling/soil treatment with a foliar application from the same mode of action group.
7. **DO NOT** re-treat a spray failure with a product from the same mode of action group.
8. Avoid using insecticide tank-mixes where both active ingredients control the same insect pests as this strategy is generally not considered best practice for resistance management.
9. Practice good crop hygiene to reduce insect/virus pressure e.g. removing severely infested plants or host weeds, and plough in old crop debris.

48

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



10. For information refer to the NSW Department of Primary Industries website: [Western flower thrips \(WFT\) insecticide resistance management plan \(nsw.gov.au\)](https://www.nsw.gov.au/western-flower-thrips-insecticide-resistance-management-plan)

Group	Chemical subgroup	Active ingredient
1A	Carbamates	methomyl
1B	Organophosphates	phorate
2B	Phenylpyrazoles (Fiproles)	fipronil
4A	Neonicotinoids	imidacloprid
5	Spinosyns	spinosad, spinetoram
6	Avermectin, Milbemycins	abamectin
8A	Alkyl halides	ethyl formate
12A + 28	Diafenthiuron + Diamides	diafenthiuron + cyantraniliprole
23	Tetronic and Tetramic acid derivatives	spirotetramat
28	Diamides	cyantraniliprole
28 + 4A	Diamides + Neonicotinoids	cyantraniliprole + thiamethoxam
29	Flonicamid	flonicamid
30	Meta-diamides, Isoxazolines	isocycloseram
UN		azadirachtin, amorphous silica
UNE	Botanical essence including synthetic, extracts and unrefined oils with unknown or uncertain MOA	orange oil
UNF	<i>Beauveria bassiana</i> strains	<i>Beauveria bassiana</i> strain PPRI 5339

Note: The table above identifies active ingredients that are either registered with the APVMA or authorised under active permits. Please consult label or permit for specific requirements.

Continued on next page.

Notes regarding the application of insecticides:

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



To ensure the most effective control of the pest:

- a) Refer to The CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides.
- b) Seek advice from the manufacturers and/or government advisory services to determine local resistance levels for particular mode of action groups.
- c) Do not exceed the maximum number of applications permitted on the insecticide label.
- d) When using insecticides to control other pests, consider the chemical group in relation to contributing to resistance development of other insect pests.
- e) When using insecticides to control insect pests consider the effect on beneficial insects and the potential to flare WFT populations.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Wide host range of plant species including Fruiting vegetables – cucurbits, Fruiting vegetables other than cucurbits, brassica vegetables, leafy vegetables, legume vegetables, root and tuber vegetables, bulb vegetables, stalk vegetables, nursery stock, ornamentals and cut flowers
Insect(s)	Serpentine leafminer (<i>Liriomyza huidobrensis</i>)

Guidelines

1. The Serpentine Leafminer is a highly polyphagous pest of around 50 different plant families including many crops and weeds. Problems with *Liriomyza* typically result from the destruction of their parasitoids by excessive use of non-selective insecticides. Therefore, an Integrated Pest Management (IPM) approach should be adopted with focus on the preservation of beneficial arthropods and monitoring of pest populations, including:
 - a. Monitoring via regular walk round of crops to determine presence of leafminers, including potential pest reservoirs in surrounding crops and weeds.
 - b. Focus on areas exposed to prevailing winds and transport routes or unloading areas.
2. *Liriomyza* leafminers are vulnerable to a wide range of generalist parasitoid and predator natural enemies, even when introduced into non-endemic regions, thus broad-spectrum products such as Organophosphates, Pyrethroids and Neonicotinoids should be avoided where possible, or used at times to minimize impact on natural enemy population, such as the end of a growing season.
3. Larval stages should be targeted by products showing systemicity, translaminar or locally systemic activity. This include Cyromazine, Abamectin, Cyantraniliprole, Chlorantraniliprole, Spinetoram, Spirotetramat (use for light infestations - suppression only).
4. Adult flies should be targeted by products with good residual and contact activity, including actives such as Abamectin, Cyantraniliprole, Chlorantraniliprole, Isocycloseram, Spinetoram.
5. When applying insecticides to this pest, key considerations should be given to:
6. Apply insecticides only when needed based on economic damage thresholds (tbd);
7. Use insecticides appropriate to the insect growth stage, e.g. systemic and translaminar acting products are required for larval stages, and contact and residual activity is key for adults.

Continued on next page.

8. Use a **medium spray quality** to ensure sufficient droplets cover the spray target to ensure the larvae ingest a lethal dose of insecticide.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



- a. Use a well calibrated, functioning boom spray with appropriate water rate for the target crop to ensure optimum spray coverage.
 - b. Use the recommended insecticide rates as stipulated on the relevant APVMA Emergency Use Permit.
 - c. Inspect the performance of the insecticide 3-4 days after application. Always document the effectiveness of each insecticide application and never re-spray a failure with an insecticide with the same mode of action. Inform the permit holder, APVMA and agronomist of any spray failures.
 - d. Resistance risk is increased where known incidences have been recorded internationally in *Liriomyza species*. This includes the following MoA groups: Carbamates (Group 1A); Cyclodienes/Organochlorines (Group 2A), Organophosphates (Group 1B); Pyrethroids (Group 3); Spinosyns (Group 5), Abamectin (Group 6) and Cyromazine (Group 17).
 - e. When rotating between modes of action, take into account the resistance management strategies for other pests which may be present.
9. When using selected insecticides targeting the serpentine leafminer, the following resistance management strategy guidelines should be implemented:
- a. If the label allows and it is required for sustained pest management, use two sequential applications of any one Mode of Action (MOA) insecticide to span a **single generation of Serpentine leafminer** (~13-26 days at 20-30°C) and then **rotate to a different MOA insecticide**;
 - b. Do not treat successive generations with products of the same MOA;
 - c. The total exposure period of any one MOA insecticide applied throughout the crop cycle (from seedling to harvest) **should not exceed 50% of the crop cycle**;
 - d. Abide by the individual label recommendation for maximum **number of allowable applications per crop per season**;
 - e. Abide by individual label recommendation for the **minimum reapplication interval** and always use the **full recommended label rates**; and
 - f. Where possible, an Area Wide Management strategy should be adopted where the same MOA insecticides are used **by all growers** in the **same time period**.

Continued on next page.

- g. As the industry learns more about how to manage this pest, this Strategy may be updated and regional-specific strategies may be developed. Check the CropLife Resistance Management [website](#) to ensure you are following the most up to date serpentine leafminer strategy.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Continued on next page.

Group*	Chemical subgroup	Example chemical (as per permit, and named crops)**	Permit
1B	Acetylcholinesterase inhibitors	Dimethoate (pulses & ornamental shrubs and trees)	PER89184 to 30.9.30
28 + 4A	Nicotinic acetylcholine receptor (NaChR) competitive modulators (Neonicotinoids) Chlorantraniliprole + thiamethoxam (Durivo®)	Thiamethoxam + Chlorantraniliprole (Brassica leafy vegetables, leafy vegetables)	PER94452 to 31.3.28
5	Nicotinic acetylcholine receptor (nAChR) allosteric modulators – Site I (Spinosyns) spinetoram (Success® Neo)	Spinetoram (snow peas, sugar snap peas and green beans)	PER87878 to 31.12.27
		Spinetoram (Ginger)	PER93550 to 31.12.26
		Spinetoram (brassica vegetables (head and leafy), cucurbits, culinary herbs, fruiting vegetables, leafy vegetables, root and tuber vegetables, stalk and stem vegetables)	PER94451 to 31.7.27
6	Glutamate-gated chloride channel (GluCl) allosteric modulators (Avermectins) abamectin (eg. Vertimec®) emamectin benzoate (eg. Proclaim® Opti) chlorantraniliprole + abamectin (Voliam® Targo)	Abamectin (Fruiting vegetables – cucurbits, fruiting vegetables other than cucurbits (excluding mushroom and sweet corn), leafy vegetables (except lettuce), legume vegetables, root and tuber vegetables, bulb onions, bulb vegetables, head cabbages, celery and rhubarb)	PER96481 to 31.7.28
		Chlorantraniliprole + abamectin (nursery stock (non-food), fruiting plants (non-bearing) cut flower and ornamentals)	PER91812 to 31.3.27
		Emamectin (suppression only: Nursery stock (non-food), fruiting plants (non-bearing) cut flower and ornamentals)	PER91812 to 31.3.27
15	Inhibitors of chitin biosynthesis affecting CHS1 (Benzoylureas eg. Dimilin®)	Diflubenzuron (nursery stock (non- food and non-bearing), cut flowers, ornamentals)	PER91812 to 31.3.27
17	Moulting disruptors , Dipteran (eg. Diptex®)	Cyromazine (broccoli, fruiting veg – cucurbits and others (excluding mushroom and corn), head lettuce, legume vegetables, root and tuber vegetables, stalk and stem vegetables, nursery stock (non-food), fruiting plants (non-bearing), cut flower, ornamentals)	PER81867 to 30.9.26
23	Inhibitors of COA Carboxylase (eg. Movento®)	Spirotetramat (suppression of snow peas, sugar snap peas, lettuce (head and leafy), parsley, green beans, celery, rhubarb, eggplant, capsicum, chilies, tomatoes.	

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Continued on next page.

Group*	Chemical subgroup	Example chemical (as per permit, and named crops)**	Permit
28	Ryanodine receptor modulators (Diamides) chlorantraniliprole (eg. Coragen®) cyantraniliprole (Benevia®) cyclaniliprole (Teppan®) chlorantraniliprole + abamectin (Voliam® Targo)	Chlorantraniliprole (spinach and silverbeet)	PER87631 to 31.3.29
		Chlorantraniliprole + Thiamethoxam (brassica leafy vegetable and leafy vegetables – seedlings)	PER94452 to 31.3.28
		Chlorantraniliprole + Abamectin (nursery stock (non-food), fruiting plants (non-bearing), cut flower, ornamentals)	PER91812 to 31.3.27
		Cyantraniliprole (celery)	PER93850 to 31.12.26
		Cyantraniliprole (bulb vegetables, fruiting vegetables, potatoes)	PER93849 to 31.12.26
		Cyclaniliprole (nursery stock (non-food), fruiting plants (non-bearing), cut flower, ornamentals)	PER91811 to 31.3.27
30	GABA-gated chloride channel allosteric modulators Isocycloseram (Simodis®)	Isocycloseram (celery, babyleaf spinach, babyleaf lettuce, kale, open leaf lettuce, parsley, coriander, shallots and leek)	PER94854 to 30.9.28
		Isocycloseram (babyleaf spinach, babyleaf chard and babyleaf lettuce)	PER94552 to 30.11.27
UN	Unknown (eg. Azamax®)	Azadirachtin (nursery stock (non-food), fruiting plants (non-bearing), cut flower, ornamentals)	PER91812 to 31.3.27

54

*Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides

**Refer to the APVMA's PubCris website (<https://portal.apvma.gov.au/permits>) to ensure permit is still active
<https://portal.apvma.gov.au/permits>

Notes

- Life cycle prediction tool developed by Cesar: <https://cesaraustralia.shinyapps.io/darabug2/>
- Hort innovation guidance document: [1303CR2 Management-guide FINAL 150620.pdf](https://www.ausveg.com.au/1303CR2_Management-guide_FINAL_150620.pdf)
([ausveg.com.au](https://www.ausveg.com.au))
- DPI factsheet:
https://keys.lucidcentral.org/keys/v3/leafminers/key/Polyphagous%20Agromyzid%20Leafminers/Media/Html/Liriomyza_huidobrensis.htm
- CABI datasheet, including information on natural enemies:
<https://www.cabi.org/isc/datasheet/30956#67C668C3-A81D-40A2-8D80-1A04AF7490B4>

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Crop(s)	Wide host range of plant species
Insect(s)	Nematodes

Guidelines

CropLife support the IRAC nematodes resistance statement which is as follows (and is available at: <https://irac-online.org/irac-nematicide-moa-classification-now-available/>):

IRAC Nematode Working Group – Nematicide Mode of Action Classification

Nematicide Resistance Risk statement

There are no substantiated examples in the scientific literature from the last century documenting cases of significant tolerance shifts or suspected resistance leading to failure of commercial agricultural nematicides against plant parasitic nematodes (PPN) under natural field conditions. Instances of these phenomena occurring have only been reported for some products under controlled laboratory conditions⁽¹⁾. Product usage approaches and nematode ecology also reduce the potential that sustained selection pressure on PPN populations occurs under field conditions. Thus overall, it can be considered that the development of resistance in PPN species to nematicides under natural field conditions is currently unconfirmed, theoretically unlikely, and poses a low risk.

The reasons underpinning this conclusion are explained below:

Unlike other plant protection products (e.g. herbicides, fungicides and insecticides), several factors limit the potential for nematicides to create high and sustained selection pressure on plant parasitic nematode (PPN) populations under field conditions:

These factors include the:

- Relatively low frequency of nematicide use in a single cropping cycle, as a proportion of the duration of the crop and the number of PPN generations. Typically, one nematicide application is made per growing season, and occasionally more in long season or perennial crops.
- Primary application methods used for nematicides in the field often target a small soil volume (e.g. crop root zone, crop beds or rows, or seed only), leaving untreated areas and host plants (weeds) that can act as refuge or source of recolonization for unexposed PPNs.
- Various nematode species have life stages (dormant or living) in host plants (e.g. crop or weeds) that may remain in the field and not be exposed to or affected by nematicide treatments. It is noteworthy to mention that very few nematicides are effective systemically in the plant against nematodes.
- Complexity of the soil environment and chemical interactions with nematicides frequently reduces product persistence, mobility and/or bioavailability, thus minimizing the likelihood of a

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



chemical product to reach a high percentage of the plant parasitic nematode population present in the field, e.g. at different soil depths or distances from the point of application, or causing exposure to multiple generations.

- Large diversity of naturally occurring organisms that may attack surviving life-stages of PPNs in soil, reducing the overall selection pressure from a single nematicide application.

Plant parasitic nematodes occur in a variety of pressures (soil population density levels) under field conditions. In some countries, and in some species, local threshold levels may be available to assess the risk of economic crop loss. Nematode management programs should be used in cases where populations of PPNs are deemed high or very high, employing multiple tactics to provide effective control and population reduction. These programs may include cultural practices e.g. crop rotations or fallow periods, solarization, nematode resistant or tolerant varieties and the application of nematicides. In cropping systems which require multiple nematicide applications within one crop cycle or on the same field over several cycles, rotation to a nematicide with a different mode of action is recommended to reduce the risk of sustained selection pressure on PPN populations.

Nematicidal products with fungicidal or insecticidal activity require additional resistance management considerations and labelling according to FRAC or IRAC guidelines.

Reduced performance of chemical nematicides can be caused by the phenomenon of Enhanced Microbial Biodegradation (EMB)⁽²⁾. This is well documented in the scientific literature and EMB should not be confused with resistance development in plant parasitic nematodes. EMB affects the level of product availability and duration of exposure of PPNs to the product, thus reducing the apparent efficacy of a nematicide application. Rotation of nematicides from different chemical classes, as well as employing other control methods such as resistant varieties and cultural methods (e.g. crop rotations) should be considered.

56

⁽¹⁾ Tolerance shifts or resistance development in PPNs under laboratory conditions:

Although few cases have been reported, continuous exposure to sub-lethal levels of a single nematicide or mode of action may lead to the development of resistant populations under laboratory conditions. This however cannot be extrapolated to field conditions.

⁽²⁾ Enhanced microbial biodegradation (EMB):

Repeated or frequent use of the same chemical nematicide in the same field soil may lead to an apparent reduction in PPN control through enhanced microbial biodegradation (EMB) of the product. EMB is the result of adaptation and increase of microbial populations that break down a particular product, therefore changing the amount of product available and/or duration of exposure of PPN's. The microbes responsible for EMB in soil may be different for different chemical classes or products, thus rotation of different nematicide types, or a reduction in the frequency of applications may decrease the likelihood of EMB occurrence.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.



Nematicides registered in Australia

Example trade name	Active	MOA
Vydate L	Oxamyl	N-1A
Nemacur	Fenamiphos	N-1B
Rugby	Cadusafos	N-1B
Counter	Terbufos	N-1B
Tervigo Nematicide	Abamectin	N-2
Indemnify Turf Nematicide	Fluopyram	N-3
Velum Prime Nematicide		
Trefinti Turf Nematicide	Cyclobutrifluram	N-3
Vaniva Tymirium Technology Nematicide		
Salibro® Rekemel	Fluazaindolizine	N-UN
Nimtz 180 EC Nematicide	Fluensulfone	N-UN
Various	Garlic extract	N-UNE
Metham Sodium / Metham Potassium	Metham	N-UNX N-UNX
Various	Dazomet (Basamid)	N-UNX
Various	Chloropicrin	N-UNX
Telone	1,3-dichloropropene	N-UNX
Various	Methyl bromide	N-UNX
Mipic	Iodomethane	N-UNX

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as of 22 July 2026. All previous versions of this strategy are now invalid.

